

Chapter

B. subtilis Probiotics for Humans, Animals and Plants: Mechanisms, Applications and Prospects

Dilan Ergün, Göksu Tartar and Ayten Yazgan-Karataş

Abstract

Bacillus bacteria have increasingly attracted research interest due to their promising probiotic properties, including strong antimicrobial, antidiarrheal, and immunostimulatory effects. They are able to stimulate the growth of natural flora, prevent intestinal inflammation, and demonstrate excellent stability even under adverse conditions. This chapter explores the probiotic applications of *Bacillus subtilis* strains in humans, animals, and plants. First, the chapter reviews the probiotic mechanisms of *B. subtilis* and the current state of research in this area. The potential applications of recombinant *B. subtilis* probiotics are also covered, along with results from preclinical studies and the future role of *B. subtilis* probiotic therapy in clinical settings. This chapter aims to survey the current landscape of *B. subtilis* probiotic research and hypothesize about its potential future impact.

Keywords: *B. subtilis*, probiotics, recombinant probiotics, biotherapy, live therapeutics

1. Introduction

The worldwide use of probiotics has increased due to their positive effects on the human microbiota and health [1]. Probiotics, as defined by the International Scientific Association for Probiotics and Prebiotics (ISAPP) panel in 2013, are live microorganisms that provide health benefits to the host when administered in sufficient quantities [2]. Most human dairy products contain probiotic bacteria, and the global market for probiotic supplements is growing significantly. *Lactobacillus* and *Bifidobacterium* are the most commonly used and commercialized genera [3, 4]. However, the genus *Lactobacillus* is microaerophilic, whereas the genus *Bifidobacterium* is strictly anaerobic. Furthermore, these genera are slow-growing and sensitive to temperature and gastric juices, leading to challenges in production, storage, and delivery processes [5]. Among these genera, the species *Bacillus subtilis* probiotics provide multifaceted advantages over conventional probiotics in several areas due to their advanced properties, including shelf stability, resistance to extreme environmental conditions, and safe interactions with humans, animals, and plants [6]. Moreover, *B. subtilis* has several distinctive characteristics, such as versatility in nutrient use for growth, spore formation, a high growth rate, growth in both aerobic

and anaerobic environments, and high enzyme and antimicrobial production, making it an attractive bacterium for probiotic research [7].

Bacillus species have been utilized as probiotics for at least 60 years, with the 1958 registration of the Italian product Enterogermina as a medicinal supplement [8]. Of the *Bacillus* species, endospore-forming soil-dwelling *B. subtilis* is the most ubiquitous bacterium in diverse ecological settings ranging from terrestrial to aquatic environments [9]. This organism's genetics and physiology have been extensively researched, and the understanding of its genetic background and physiology is second only to that of *Escherichia coli* [10, 11]. Recent probiotic studies of *B. subtilis* have made significant advances in clarifying the spectrum of activity that makes this bacterium one of the most appealing probiotics [5]. *B. subtilis* strains have been studied as probiotics for their antibacterial, antifungal, antiviral, antidiarrheal, and immunostimulatory effects, as well as for their potential to compete with pathogens for survival, inhibit intestinal inflammation, and stimulate the growth of normal intestinal flora [12–17]. This chapter discusses the applications of *B. subtilis* probiotics in humans, animals, and plants. First, the chapter reviews the probiotic mechanisms of *B. subtilis* and the current state of research in this area. The potential applications of recombinant *B. subtilis* probiotics are also covered, along with results from preclinical studies and the future role of *B. subtilis* probiotic therapy in clinical settings.

2. Mechanisms of action of *B. subtilis* probiotics

B. subtilis' probiotic effects on the host can be attributed to a variety of mechanisms, including antimicrobial effect, antidiarrheal effect, immunostimulatory effect, competitive exclusion of pathogens, prevention of intestinal inflammation, and stimulation of growth of intestinal normal flora [18].

2.1 Synthesis of antimicrobial agents

Both industry and healthcare have made extensive use of *Bacillus* spp. due to their ability to produce a wide range of biologically active metabolites, particularly antimicrobials [19]. Pathogenic infections are a leading cause of morbidity and mortality worldwide, and while synthetic antibiotics are used to control pathogenic overgrowth, their indiscriminate use leads to the emergence of multidrug-resistant pathogenic bacteria. *B. subtilis*, one of the most prominent species, has shown promise in this area by serving as a viable therapeutic antimicrobial, as it synthesizes 66 antibiotics and allocates 4–5% of its genome to antibiotic biosynthesis [1, 7, 20]. The antibiotics synthesized by *B. subtilis* have diverse structures and antimicrobial properties. Specifically, *B. subtilis* synthesizes antimicrobial agents that disrupt cell walls and plasma membranes and interfere with intracellular processes [19]. Bacilysin exerts its effect by targeting the cell walls of *Candida albicans* and foodborne pathogens such as *Escherichia coli*, *Salmonella enterica* and *Bacillus cereus* [21, 22]. *B. subtilis* biosynthesizes lipopeptide antibiotics such as surfactin and fengicyn, which exert their antimicrobial activity against harmful microorganisms such as plant pathogens such as *Rhizopus stolonifer* through their interaction mechanisms with phospholipid bilayer membranes [23, 24]. Some agents inhibit pathogens by disrupting intracellular processes without disrupting cell walls and plasma membranes; for example, sublancin, a glycosylated peptide produced by *B. subtilis* 168, shows antibacterial activity

against methicillin-resistant *Staphylococcus aureus* by disrupting the central dogma [25, 26]. In this regard, *B. subtilis* probiotics are promising antibiotic substitutes with remarkable antimicrobial properties, highlighting their potential as a novel strategy to overcome microbial infections [27]. For example, supplementation with *B. subtilis* PB6 in *E. coli*-infected broilers reduced the mortality rate from 14% to 8%, while antibiotic administration reduced it to 6% [28]. Similarly, compared to negative controls, supplementation with *B. subtilis* DSM 32315 and Nasarin reduced mortality by 6.5% and 9.1%, respectively [29]. *B. subtilis* has also been shown to reduce *Salmonella typhimurium* infection in cell culture studies, in addition to mucosa-associated lymphoid tissue (MALT) healing [30].

2.2 Immunostimulatory effect

B. subtilis stimulates both specific and non-specific immunity, thereby improving protection against pathogens. *Bacillus* spores have been shown to stimulate the immune system by oral administration in numerous studies in humans and animal models. *B. subtilis* spores induce specific humoral and cell-mediated immune responses. The development of both innate and adaptive immune responses in the host is significantly influenced by the interaction between *B. subtilis* spores and macrophages [5]. Studies on macrophages and *B. subtilis* spore co-cultures showed that *B. subtilis* significantly increased the secretion of pro-inflammatory cytokines, including tumor necrosis factor-alpha (TNF- α), interferon-gamma (IFN- γ), interleukin-1 beta (IL-1 β), IL-6, IL-8 and IL-12 [31–33]. Lefevre et al. [34] demonstrated that supplementation of elderly humans with *B. subtilis* CU1 significantly increased fecal and salivary secretory IgA concentrations compared to placebo. Administration of *B. subtilis* MBTU PBBM1 spores to mouse models resulted in T lymphocyte proliferation and accumulation of IgG and IgA antibodies, indicating the potential of *B. subtilis* MBTU PBBM1 spores to enhance both humoral and cellular immunity in mice [35]. Bai et al. [36] showed that administration of *B. subtilis* fmbJ to broilers significantly increased serum IgG and IgA levels. In addition, a study by Li et al. [37] in growing-finishing pigs showed that *B. subtilis* increased serum IgG levels, suggesting that supplementation with *B. subtilis* significantly stimulates the immune response. A study in a rabbit model showed that the combination of *Bacteroides fragilis* and *B. subtilis* had a positive effect on the development of gut-associated lymphoid tissue (GALT) and the diversification of the pre-immune antibody repertoire. Furthermore, two stress responses controlled by Spo0A, including sporulation and TapA (formerly YqxM) secretion, were found to be necessary for GALT development in *B. subtilis* [38].

2.3 Maintenance of intestinal homeostasis

Probiotic *B. subtilis* strains have shown promising results in supporting gastrointestinal health and exhibiting anti-inflammatory properties [5, 6]. Patch et al. [39] found that *B. subtilis* supplementation significantly improved symptoms of constipation, indigestion and dyspepsia compared to the placebo control group. Further research has shown that stool quality was significantly improved following *B. subtilis* supplementation, while stool frequency per day and abdominal noises were significantly reduced [40, 41]. In addition, anti-inflammatory immune cell populations increased in response to *ex vivo* lipopolysaccharide stimulation of peripheral blood mononuclear cells [42]. Studies in piglets and growing-finishing

pigs showed that *B. subtilis* supplementation improved the efficiency of digestion and metabolism of nutrients and had antioxidant effects. *B. subtilis* was shown to have a positive effect on intestinal histomorphology and its supplementation could promote a more favorable mucosal structure and larger absorptive areas of the luminal villi, thus promoting better intestinal development [37, 43, 44]. This finding suggested the reason for the improved digestive efficiency [37]. It was also reported that administration of *B. subtilis* to pigs increased serum superoxide dismutase levels, and after administration of *B. subtilis* fmbJ, reactive oxygen species (ROS) levels in liver mitochondria of broilers were significantly decreased, and the expression of antioxidant enzyme genes including nuclear factor erythroid 2-related factor 2 (Nrf2), haem oxygenase 1 (HO-1), superoxide dismutase (SOD), and glutathione peroxidase (GPx) were improved. Irritable bowel syndrome and inflammatory bowel disease (IBD), which includes Crohn's disease and ulcerative colitis, have been shown to be associated with oxidative stress and may benefit from antioxidant effects that help maintain and restore intestinal function [45]. Studies in broilers showed that supplementation with *B. subtilis* increased goblet cells in the intestinal tract by 19% and reduced the severity of inflammatory cell infiltration in the jejunum [46, 47]. Research in mice showed that histological damage caused by IBD was reduced by the ingestion of *B. subtilis*, which upregulated the expression of tight junction proteins and suppressed cytokine expression to mitigate intestinal epithelial damage [48]. Furthermore, competence and sporulation factor (CSF) synthesized by *B. subtilis* was found to activate the p38 MAP kinase and protein kinase B (Akt) survival pathways in intestinal epithelial cells, preventing oxidant-induced intestinal epithelial cell injury and loss of barrier function after its OCTN2-mediated transport [49]. CSF also reduced intestinal inflammation and improved survival in mice with lethal colitis [50].

2.4 Maintenance of gut microbiota

B. subtilis administration has been shown to influence the gut microbiota profile by stimulating the growth of normal gut flora and by competitively eliminating pathogens [51]. Studies conducted in broilers showed that *B. subtilis* promoted the presence of beneficial bacteria in the gut such as *Enterococcus*, *Lactobacillus* spp., *Bifidobacterium* spp., *Firmicutes*, *Christensenellaceae*, *Caulobacteraceae*, *Ruminococcus*, *Lachnospirillum*, and *Anaerostipes* while reducing *Vampirovibrio*, *Escherichia/Shigella*, *Parabacteroides*, *Clostridium*, *Coliforms*, *Campylobacter*, *Bacteroidetes*, and *E. coli* [52–58]. It has also been shown that adherence of *Salmonella enteritidis*, *Listeria monocytogenes*, and *Escherichia coli* pathogens to the HT-29 cell line was inhibited by *B. subtilis* P223 [16]. Furthermore, the *B. subtilis*-treated model organism *Bombyx mori* L. had a higher abundance of *Enterococcus* compared to the control group [59]. Studies in pigs and piglets exhibited that *B. subtilis* supplementation increased the numbers of *Lactobacilli*, *Leucobacter* and *Cupriavidus*, *Coproccoccus*, and *Bifidobacterium*, as well as colonic *Succiniclasticum* cells the numbers of *Clostridium perfringens* and *E. coli* [43, 54, 60, 61]. Other studies have demonstrated that *B. subtilis* favors the intestinal colonization of some beneficial bacteria, including *Lactobacillus*, *Leucobacter*, *Bifidobacteria*, *Megasphaera*, *Coproccoccus*, *Gemmatimonadete*, *Acidobacteria*, *Ruminococcaceae* UCG-013 cc, and *Prevotella* [43, 62, 63]. Administration of *B. subtilis* has been shown to alter the microbial composition and optimize the structure of the gut microbiome, contributing to its beneficial effects as a probiotic [37, 43, 44, 63].

3. Applications of probiotic *B. subtilis*

3.1 For humans

Human studies have demonstrated the potential of *B. subtilis* probiotics to effectively treat various intestinal infections and diseases, including antibiotic-associated diarrhea, ulcerative colitis, acute intestinal infections, irritable bowel syndrome, and other infectious pathologies [5, 6]. In addition, probiotic *B. subtilis* has been shown to be effective in the treatment of immunodeficiency [64], Alzheimer's disease [65] and Parkinson's disease [66]. *B. subtilis*-derived probiotics have been shown to increase human longevity by slowing the progression of age-related diseases [67]. In addition, *B. subtilis* supplementation has been beneficial in reducing risk factors associated with cardiovascular disease by lowering cholesterol levels in the human bloodstream [68, 69]. Furthermore, the use of *B. subtilis* natto DG101 in conjunction with appropriate medication has become a promising and innovative intervention in the fight against the global diabetes epidemic [70].

B. subtilis has also shown promise as an adjunctive treatment for *Helicobacter pylori* eradication therapy and dysbacteriosis in neonates and children [5, 6]. Given its ability to favorably alter the gastrointestinal microflora, *B. subtilis* probiotics are being considered as a daily supplement to support overall gut and digestive health in individuals. Probiotic strains of *B. subtilis* have been shown to have beneficial effects on GI health, including improvements in constipation [39], GI quality of life [71], body composition [72–74], stool regularity [40, 41], GI discomfort [75], bloating and belching [76], and microbiome composition [41, 77]. Additional benefits have been observed in glucose metabolism [73], amino acid, purine and vitamin B metabolism [59], GI hormonal response [68, 78], anti-inflammatory activity [42], lipid metabolism [68, 72], endothelial function [68], bone density [79] and immune response [34].

In recent years, engineered probiotics have gained considerable attention as a live therapy to maintain healthy gut microbiota and treat microbial infections. *Lactobacillus rhamnosus* GG (LGG) is a widely studied probiotic for use in ulcerative colitis [80], diarrhea [81] and atopic dermatitis [82]. A major secreted protein, p75, enhances its probiotic activity by maintaining intestinal epithelial cell proliferation and survival by stimulating activation of Akt, suppressing cytokine-induced epithelial cell apoptosis, increasing cell proliferation and reducing tumor necrosis factor (TNF)-induced colonic epithelial damage [83, 84]. However, maintaining the viability of these probiotics during food processing and in the gut remains a challenge [85, 86]. A recent study anchored the p75 protein from *L. rhamnosus* GG to *B. subtilis* via CotG, demonstrating its antibacterial activity against the foodborne pathogen *Listeria monocytogenes* [27]. Researchers have also investigated *B. subtilis* probiotics as a vaccine delivery system. These spore probiotics enhance immune responses by presenting antigens in the gut [87]. For example, researchers have proposed *B. subtilis* spores expressing *H. pylori* urease B (UreB) protein on their surface as a promising oral vaccine against *H. pylori* infection. In experimental mouse models, oral administration of UreB-expressing recombinant *B. subtilis* spores resulted in a significant humoral immune response. Furthermore, oral administration of these engineered spores reduced *H. pylori* load in infected mice by 84% compared to the control group [88]. Several studies have highlighted the importance of surface display of antigens on *B. subtilis* probiotics, such as the C fragment of tetanus toxin (TTFC) [89], the tegument protein of *Clonorchis sinensis* [90] and the SARS-CoV-2 spike protein [91]. These

studies have shown that oral administration of these engineered *B. subtilis* probiotics induces robust immune responses.

Researchers are exploring novel approaches to promote gut health, modify metabolic pathways and improve overall health through the development of recombinant *B. subtilis* probiotic candidates, paving the way for a new era in the treatment of infectious diseases and metabolic disorders. For example, a recent study evaluated the efficacy of a novel recombinant *B. subtilis* SP-64 strain expressing human alcohol dehydrogenase-4 (ADH4) in mice to treat diet-induced metabolic disorders such as obesity, fatty liver disease and type 2 diabetes [1]. Long-term ethanol production in the gastrointestinal (GI) system can damage the gut microbiota and cause organ dysfunction. It is necessary to reduce ethanol exposure to internal organs and increase ADH4 activity, which is necessary for alcohol metabolism [92]. In the study, increased rates of ethanol removal were achieved with the customized *B. subtilis*. Treatment with this customized probiotic significantly reduced weight gain in mice, promoted glucose utilization and prevented pancreatic damage from a high-fat diet [1]. A recent study investigated a lactate oxidase-expressing recombinant *B. subtilis* as a therapeutic probiotic candidate for metabolic disorders associated with elevated blood lactate levels [93]. Anaerobic glycolysis can lead to elevated lactate levels associated with diseases such as cancer, diabetes and mitochondrial dysfunction [94–96]. Traditional techniques for treating elevated lactate levels, such as small molecule inhibitors or enzyme replacement, have limited clinical efficacy [97–99]. In this study, lactate oxidase (LOX)-expressing recombinant *B. subtilis* probiotic candidates were orally administered to the mouse models to improve the rate of removal of elevated blood lactate levels and combat associated metabolic diseases. Oral administration of these LOX-expressing recombinant spore-based *B. subtilis* probiotics significantly reduced elevated blood lactate levels in the animal models without affecting the immune system, liver or gut microbiome [93].

3.2 For animals

Livestock production has expanded significantly to meet the needs of a growing global population, with antibiotics being used as growth promoters to improve feed conversion rates [100–102]. However, this has led to the development of antibiotic-resistant pathogenic bacteria that pose a threat to human health and the economy [103, 104]. The rise of antimicrobial resistance has reduced effective treatments, reduced animal productivity and increased the risk of infectious diseases [105]. The use of antibiotic growth promoters as feed additives has been prohibited in many countries due to regulatory pressure and consumer demand [106]. Alternatives such as probiotics, feed enzymes, prebiotic oligosaccharides, synbiotics, essential oils and immunostimulants have gained interest to maintain current animal production metrics [107]. As a promising probiotic alternative, *B. subtilis* strains have long been used as feed additives to promote animal growth and prevent disease [108]. Its ability to be produced in the form of endospores allows it to rapidly reactivate in the upper intestine to secrete proteases, facilitating the breakdown of complex carbohydrates in plant diets [109–111]. *B. subtilis* produces antimicrobial agents that fight intestinal pathogens and improve intestinal health and food absorption [19–24, 47, 51].

Animal feed production processes typically require high temperatures, and spore-producing bacteria are well suited to these conditions. In the production of functional animal feeds, the ability to sporulate, grow under both aerobic and anaerobic conditions, and survive inside and outside of humans and animals makes the *B. subtilis*

strains one of the most versatile and attractive probiotic bacteria [7, 107]. Several studies have evaluated the effects of *B. subtilis* probiotics on broiler growth performance compared to antibiotics. Lee et al. [112] found that broilers supplemented with *B. subtilis* had higher body weights than those supplemented with 60 mg/kg salinomycin. Teo and Tan [28] showed that the administration of *B. subtilis* PB6 to *E. coli*-infected broilers improved the average daily gain from day 21 to 41 by 7.5% compared to the administration of a combination of 16.7 mg/kg bacitracin zinc and 3.3 mg/kg colistin sulfate. In addition, recombinant *B. subtilis* probiotics in *Eimeria maxima*-infected broilers significantly improved gut integrity, enhanced innate immunity, and improved growth performance in young chickens [113]. In addition, a study proposed *Clostridium perfringens* alpha toxoid expressing recombinant *B. subtilis* as a promising vaccine for the prevention of necrotic enteritis in poultry [114]. *B. subtilis* spore probiotics expressing a truncated Newcastle disease virus haemagglutinin-neuraminidase protein and a *B. subtilis* strain expressing the duck hepatitis A virus type 1 VP1 protein significantly induced strong systemic and mucosal immune responses in mice, highlighting the potential of *B. subtilis* probiotics as an immunization agent for poultry [115, 116]. The *B. subtilis* probiotic strain was also genetically modified to detoxify poultry feed by inserting the dioxin degradation system of *Sphingomonas wittichii*. It significantly reduced dioxin levels, demonstrating its potential as a feed additive to detoxify poultry diets and provide safe consumption for chickens [117].

Newborn piglets have a sterile gut that eventually develops their characteristic microbiome through interaction with the sow and surrounding environment. The periods immediately following birth and post-weaning represent the most vulnerable stages in a pig's lifecycle, where the elimination of antimicrobial substances from feed could impact mortality rates [118, 119]. A recombinant *B. subtilis* probiotic expressing antimicrobial peptide was developed to address these issues in swine production. Tests on the constructed recombinant probiotics *in vitro* exhibited antimicrobial activity against *E. coli*, *S. typhimurium*, *Haemophilus parasuis*, and *S. aureus*. Furthermore, administration of these engineered probiotics in weaned piglets significantly enhanced the growth and reduced the incidence of diarrhea from enteropathogenic *E. coli* challenge compared to non-supplemented piglets [120].

Porcine epidemic diarrhea virus (PEDV) is a serious pig disease causing high mortality from diarrhea in piglets [121–123]. Currently, available PEDV vaccines lack efficacy, resulting in inadequate treatment [124, 125]. As *B. subtilis* probiotics are known to improve immune regulation and promote gut health in pigs when administered orally [126], engineering *B. subtilis* probiotics to provide immunization against PEDV infection is a promising approach to overcome the current challenges in treating PEDV infection [127]. These probiotics could also be developed as feed additives with enhanced nutritional value. Previous work has shown that tryptophan-excreting *B. subtilis* probiotics can produce tryptophan *in situ* to supplement pig diets, offering the potential benefits of reducing environmental pollution from nitrogen sources and feed costs [128].

Live microbial feed additives, such as probiotics, may not only alter the rumen microbiota but also induce strong host immune responses [129, 130]. Higher feed efficiency and improved milk production have been demonstrated in cows fed *B. subtilis* natto fermentation products, suggesting that *B. subtilis* natto alters rumen fermentation patterns [129, 131]. *Haemonchus contortus* is an important gastrointestinal parasite that infects sheep, causing blood loss and anemia. *H. contortus* infection affects the gastrointestinal microbial population, inhibiting protein digestion and growth performance in infected animals [132, 133]. A recent study showed that HcGAPDH

protein from *H. contortus* expressing a recombinant probiotic *B. subtilis* strain induced immune responses in sheep and reduced worm burden [134].

3.3 For plants

Biological products containing live bacterial cells and/or their metabolites have great potential as next-generation methods of crop protection. These technological advances now allow plant breeders to potentially avoid reliance on chemical pesticide and fungicide applications in some cases. *B. subtilis*, in particular, has shown promise as an effective biological control alternative to chemical control [135, 136]. *B. subtilis* is a common plant growth-promoting rhizobacterium (PGPR) that enhances plant development and yield. It protects the plant against stress through biofilm development, lipopeptide production, siderophore secretion and exopolysaccharide release. It acts as an effective denitrifying agent in the agroecosystem and maintains soil health through environmentally sustainable remediation approaches [137]. For example, the use of *B. subtilis* as a biofertilizer in fertigated Tarocco blood orange groves has been shown to improve fruit quality, soil fertility, growth and nutrient uptake, and to reduce the need for chemical fertilizers [138]. Field trials using *B. subtilis* CPA-8 have reduced *Monilinia* spp. plant infections, demonstrating its potential as an environmentally friendly alternative to fungicides [135]. The chitinase from *Bacillus pumilus* was expressed on *B. subtilis* spores and its biopesticidal activity was demonstrated against fungal pathogens such as *Rhizoctonia solani* and *Trichoderma harzianum*, which are major threats to maintaining sustainable agriculture, highlighting the potential of these novel biocontrol agents [139].

4. Clinical trials

Functional gastrointestinal disorders affect 25–40% of the world's population. They are chronic conditions with persistent symptoms, including functional bowel disorders, irritable bowel syndrome, constipation, diarrhea and bloating [39, 140]. Antibiotics are often ineffective or contraindicated for long-term use, whereas probiotics offer health benefits without resistance [141]. A clinical trial investigated the use of probiotic *B. subtilis* (BG01-4TM) in the treatment of functional gastrointestinal disorders and found that BG01-4TM relieved constipation symptoms, suggesting its potential in the treatment of GI dysfunction [39]. A study of 400 adult patients with moderate-to-severe diarrhea-predominant irritable bowel syndrome (IBS-D) found that *B. subtilis* containing Bio-Kult®, a multi-strain probiotic, reduced abdominal pain and improved bowel movements, suggesting potential benefits for IBS-D patients [142]. A clinical trial in 76 healthy subjects found that daily *B. subtilis* BS50 supplementation for 6 weeks improved bloating, belching and flatulence, suggesting a safe approach to gastrointestinal discomfort [76]. A study in 81 healthy adults aged 18–50 years evaluated the tolerability and safety of *B. subtilis* R0179. The study found no adverse effects on general well-being or gastrointestinal function. *B. subtilis* R0179 was well tolerated at doses ranging from 0.1 to 10×10^9 cfu/day [75].

In a recent study, two cases of type 2 diabetes were well controlled with *B. subtilis* natto DG101 probiotics. Probiotics helped control blood glucose and insulin levels in two cases of type 2 diabetes in obese patients. This study shows the potential of *B. subtilis* probiotics in managing type 2 diabetes and in maintaining health and managing disease [70].

5. Future prospects

Recently, *B. subtilis* as a probiotic has gained attention for its applications in skin care. A study in mice showed that a *B. subtilis*-supplemented Pluronic F-127 hydrogel could modulate the skin microbiome, revealing its potential in the treatment of skin dysbiosis-related diseases such as atopic dermatitis, acne, rosacea, and psoriasis [143]. Furthermore, topical application of *B. subtilis*-substituted polyvinyl alcohol (PVA) microparticles effectively reduced the healing time of open wounds in mice, indicating the potential of *B. subtilis* as an effective therapeutic agent for acute or chronic open wounds [144]. Despite its promising potential, there is a lack of scientifically validated clinical data on the efficacy and safety of probiotic *B. subtilis* formulations for the treatment of skin disorders, which could be urgently pursued as future research.

Psychobiotic strains influence central nervous system (CNS)-related functions and behaviors mediated by the gut-brain axis (GBA) through a number of different pathways [145]. There is growing interest among researchers in the field of microbial-based psychopharmaceuticals. An imbalance in the gut microbiota has been linked to a range of health problems, including inflammatory bowel disease, obesity, type 2 diabetes, and even mental health conditions such as autism, Alzheimer's disease (AD), depression and anxiety [146]. In recent years, preclinical and some clinical studies of psychobiotics have yielded encouraging results in the treatment of neurodegenerative and depressive disorders [65, 147]. Bacteria of the genera *Lactobacillus* and *Bifidobacterium* have been most extensively studied for their psychobiotic effects [146, 147]. Conversely, recent studies have shown that *B. subtilis* strains also have promising psychobiotic properties. For example, studies using chickens as a model organism have shown that *B. subtilis* has beneficial effects on growth performance, intestinal microstructure and digestive enzymes, synthesis and release of antimicrobial and antibiotic compounds, immunity and neurotransmitter activities, including serotonin, and animal behavior following various stressors. In addition, *B. subtilis* has the ability to overproduce L-tryptophan, which subsequently increases 5-HT in the hypothalamus, acting as an antidepressant and anti-anxiety agent, relieving nervous tension in mice and reducing aggressive behavior in laying hens [148]. Besides serotonin, *B. subtilis* has the ability to produce gamma-aminobutyric acid (GABA) [149, 150]. GABA is one of the most important inhibitors in the brain, regulating many physiological and psychological processes. Its dysfunction has been implicated in anxiety and depression. It also plays an important role in the prevention of neurological diseases, cancer, type 1 diabetes and immune disorders [151]. Nitric oxide (NO), another neurotransmitter involved in several GI functions, including maintaining vascular tone, modulating immune responses, transmitting nerve impulses, and facilitating GI motility by relaxing intestinal muscles, could be produced by *B. subtilis* [152]. Alzheimer's disease (AD) is a major cause of dementia among diseases of the nervous system. The exact etiology of Alzheimer's disease remains incompletely understood. There is no definitive and effective treatment for the disease. *B. subtilis* NCIB 3610 has been extensively studied for its ability to form well-structured, three-dimensional biofilms *in vitro*, which is one of the many potential pathways by which it exerts a probiotic effect [153]. It has been shown to delay the onset of age-related neurodegenerative and cognitive deficits in a *Caenorhabditis elegans* model of AD [65]. The anti-AD effects of *B. subtilis* are directly related to the ability of the probiotic to synthesize a quorum-sensing peptide—Competence Sporulation Factor (CSF)—and to gut-associated biofilm development. Alpha-synuclein (α Syn), a highly soluble

unfolded protein, has been implicated in Alzheimer's disease and Parkinson's disease. A reduction in DA levels in α Syn has been observed in post-mortem AD brains, which is associated with cognitive decline. Another *B. subtilis* PXN21 has been observed to inhibit, delay and even reverse α Syn aggregation in young and old *C. elegans* models by altering the sphingolipid metabolic pathway. Despite these promising results, scientifically validated research into their psychobiotic effects has been limited. Their effects may vary depending on factors such as strain type, dosage and individual differences. Comprehensive molecular mechanism studies, animal experiments and clinical trials are therefore essential to refine their use [65, 154, 155].

The potential therapeutic efficacy of *B. subtilis* against coronavirus disease (COVID-19) has recently been elucidated [156]. By modulating the immune system, *B. subtilis* appears to help prevent excessive inflammation and thus improve patient outcomes. Preliminary evidence suggests that this bacterium may have an inhibitory effect on viral replication. This observation warrants further investigation, particularly given the lack of clarity regarding its specific activity against the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus. *B. subtilis* is still being tested as a treatment for patients with COVID-19 [156] and further studies are needed to confirm its potential and understand how it works.

Recent advances in “omics” technologies, including genomics, transcriptomics, proteomics and metabolomics, have significantly improved our understanding of *B. subtilis* at the molecular level. By integrating omic approaches with systems biology studies, researchers have gained insight into the networks and pathways that control the physiology of this model bacterium [157]. Advances in synthetic biology and genome editing techniques such as CRISPR-Cas9 have led to the development of engineered probiotics for various health problems. These probiotics can be used to treat inflammatory bowel disease, cancer and neurological disorders. *B. subtilis*, known for its non-pathogenicity, excellent protein secretion ability, adaptive metabolism and resistance, makes it an ideal candidate for metabolic engineering and synthetic biology to produce live therapeutics for disease treatment [6, 158]. An intriguing example of such applications is the genetically engineered probiotic *Bacillus subtilis* ZB183, which is Food and Drug Administration (FDA)-compliant and has been developed with the ability to reduce alcohol hangover. This product has been marketed by ZBiotics since 2019 [159, 160]. In addition, in 2024, ZBiotics commercialized another engineered probiotic bacterium, ZB423™, to convert sugar into fiber continuously throughout the day. This unique probiotic produces levansucrase, an enzyme that splits sugar into glucose and fructose, and redirects the fructose to form a rare fiber called levan. This enzyme is delivered directly to the gut, maximizing the benefits and delivering the fiber in a way that is preferred by the microbiome [161, 162]. These two recombinant commercial examples represent a significant advance demonstrating the success and safety of engineered *B. subtilis* probiotics, the number of which is expected to increase rapidly in the near future.

6. Conclusions

The *Bacillus* genus of bacteria has recently become the subject of considerable research interest, due to the significant potential health benefits they offer to humans, animals and plants. For humans, *B. subtilis* offers a wide range of health-promoting properties, including improved digestion, stimulation of the growth of natural microflora, circulatory blood homeostasis, stimulation of the immune system, prevention

of intestinal inflammation, and potential protection against pathogenic invaders. Clinical trials have demonstrated the safety and efficacy of *B. subtilis* as a dietary supplement for the treatment of diseases affecting the gastrointestinal (GI) tract and type-II diabetes. In addition, preclinical studies have also yielded promising results for the management of cardiovascular diseases, inflammatory disorders, dermatological diseases, neurodegenerative and depressive conditions, and even SARS-CoV-2 infection.

B. subtilis is being used more in livestock farming because it improves feed efficiency and eliminates the need for antibiotics. This is achieved by improving nutrient digestibility, reducing pathogen load, strengthening mucosal physiology and improving gut immunity and microbiota diversity in various livestock species. Together with these beneficial effects, the capacity of *B. subtilis* to form spores at high temperatures and to survive in low pH environments renders it a robust strain that can be developed into an in-feed probiotic supplement.

The *B. subtilis* species is of great importance in maintaining plant health, promoting plant growth and fighting against pathogens. *B. subtilis* has been demonstrated to be highly effective in combating pathogens, enhancing nutrient availability and enhancing stress tolerance in plants. This highlights its significant potential in shaping the trajectory of sustainable agriculture.

The advent of synthetic biology with the advancement of genome editing techniques such as CRISPR-Cas9 has engendered promising prospects for the customization of *B. subtilis* strains. As a proficient chassis cell, these organisms can be modified to alter their intrinsic properties, thereby facilitating the development of novel solutions for the treatment of a spectrum of human, animal and plant diseases. Given its status as a commensal organism present in humans, animals and plants, its exceptional resilience even in adverse conditions and its designation as a Generally Recognized as Safe (GRAS) organism, engineered *B. subtilis* probiotics are poised to assume an even more pivotal role in sustainable practices and innovative solutions to the challenges of human, animal and plant health in the near future.

Author details

Dilan Ergün^{1,2}, Göksu Tartar^{1,2} and Ayten Yazgan-Karataş^{1,2*}

1 Department of Molecular Biology and Genetics, Faculty of Science and Letters, Istanbul Technical University, Istanbul, Türkiye

2 Dr. Orhan Ocalgiray Molecular Biology Biotechnology and Genetics Research Center, Istanbul Technical University, Istanbul, Türkiye

*Address all correspondence to: karatasay@itu.edu.tr

IntechOpen

© 2024 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Singh RP, Shadan A, Ma Y. Biotechnological applications of probiotics: A multifarious weapon to disease and metabolic abnormality. *Probiotics and Antimicrobial Proteins*. 2022;**14**(6):1184. DOI: 10.1007/S12602-022-09992-8
- [2] Hill C, Guarner F, Reid G, Gibson GR, Merenstein DJ, Pot B, et al. Expert consensus document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nature Reviews Gastroenterology & Hepatology*. 2014;**11**(8):506-514. DOI: 10.1038/NRGASTRO.2014.66
- [3] Chen J, Chen X, Ho CL. Recent development of probiotic bifidobacteria for treating human diseases. *Frontiers in Bioengineering and Biotechnology*. 2021;**9**:770248. DOI: 10.3389/FBIOE.2021.770248/
- [4] Venkatesh GP, Kuruvalli G, Syed K, Reddy VD. An updated review on probiotic production and applications. *Gastroenterology Insights*. 2024;**1**:221-236. DOI: 10.3390/GASTROENT15010016
- [5] Suva M, Sureja V, Kheni D. Novel insight on probiotic *B. subtilis*: Mechanism of action and clinical applications. *Journal of Current Research in Scientific Medicine*. 2016;**2**(2):65. DOI: 10.4103/2455-3069.198381
- [6] Williams N, Weir TL. Spore-based probiotic *B. subtilis*: Current applications in humans and future. *Perspectives*. 2024;**10**(2):78. DOI: 10.3390/FERMENTATION10020078
- [7] Olmos J, Paniagua-Michel J. *B. subtilis* a potential probiotic bacterium to formulate functional feeds for aquaculture. *Journal of Microbial & Biochemical Technology*. 2014;**06**(07):361-365. DOI: 10.4172/1948-5948.1000169
- [8] Cutting SM. *Bacillus* probiotics. *Food Microbiology*. 2011;**28**(2):214-220. DOI: 10.1016/J.FM.2010.03.007
- [9] Earl AM, Losick R, Kolter R. Ecology and genomics of *B. subtilis*. *Trends in Microbiology*. 2008;**16**(6):269-275. DOI: 10.1016/J.TIM.2008.03.004
- [10] Kunst F, Ogasawara N, Moszer I, Albertini AM, Alloni G, Azevedo V, et al. The complete genome sequence of the gram-positive bacterium *B. subtilis*. *Nature*. 1997;**390**(6657):249-256. DOI: 10.1038/36786
- [11] Driks A. Proteins of the spore core and coat. In: Sonenshein H, Losi, editors. *B. subtilis* and Its Closest Relatives. Washington, D.C: ASM Press; 2014. pp. 527-535. DOI: 10.1128/9781555817992.ch36
- [12] Ritter A, Correa A, Veras F, Brandelli A. Characterization of *B. subtilis* available as probiotic. *Journal of Microbiology Research*. 2018;**8**(2):23-32. DOI: 10.5923/j.microbiology.20180802.01
- [13] Starosila D, Rybalko S, Varbanetz L, Ivanskaya N, Sorokulova I. Anti-influenza activity of a *B. subtilis* probiotic strain. *Antimicrobial Agents and Chemotherapy*. 2017;**61**(7):e00539-17. DOI: 10.1128/AAC.00539-17
- [14] Urdaci MC, Lefevre M, Lafforgue G, Cartier C, Rodriguez B, Fioramonti J. Antidiarrheal action of *B. subtilis* CU1 CNCM I-2745 and *L. plantarum* CNCM I-4547 in Mice. *Frontiers in Microbiology*. 2018;**9**:1537. DOI: 10.3389/FMICB.2018.01537

- [15] Huang JM, la Ragione RM, Nunez A, Cutting SM. Imm-unostimulatory activity of *Bacillus* spores. *FEMS Immunology and Medical Microbiology*. 2008;**53**(2):195-203. DOI: 10.1111/J.1574-695X.2008.00415.X
- [16] Jeon HL, Lee NK, Yang SJ, Kim WS, Paik HD. Probiotic characterization of *B. subtilis* P223 isolated from kimchi. *Food Science and Biotechnology*. 2017;**26**(6):1641-1648. DOI: 10.1007/S10068-017-0148-5
- [17] Rhayat L, Maresca M, Nicoletti C, Perrier J, Brinch KS, Christian S, et al. Effect of *B. subtilis* strains on intestinal barrier function and inflammatory response. *Frontiers in Immunology*. 2019;**10**:564. DOI: 10.3389/FIMMU.2019.00564
- [18] Leser TD, Knarreborg A, Worm J. Germination and outgrowth of *B. subtilis* and *B. licheniformis* spores in the gastrointestinal tract of pigs. *Journal of Applied Microbiology*. 2008;**104**(4):1025-1033. DOI: 10.1111/J.1365-2672.2007.03633.X
- [19] Tran C, Cock IE, Chen X, Feng Y. Antimicrobial *Bacillus*: Metabolites and their mode of action. *Antibiotics*. 2022;**11**(1):88. DOI: 10.3390/ANTIBIOTICS11010088/S1
- [20] Sorokulova I. Modern status and perspectives of *Bacillus* bacteria as probiotics. *Journal of Probiotics and Health*. 2013;**1**(4):e106. DOI: 10.4172/2329-8901.1000e106
- [21] Wang T, Liu X-h, Wu M-b, Ge S. Molecular insights into the antifungal mechanism of bacilysin. *Journal of Molecular Modeling*. 2018;**24**(5):112. DOI: 10.1007/S00894-018-3645-4
- [22] Nannan C, Vu HQ, Gillis A, Caulier S, Nguyen TTT, Mahillon J. Bacilysin within the *B. subtilis* group: Gene prevalence versus antagonistic activity against Gram-negative foodborne pathogens. *Journal of Biotechnology*. 2021;**327**:28-35. DOI: 10.1016/J.JBIOTECH.2020.12.017
- [23] Tao Y, Bie X-m, Lv F-x, Zhao H-z, Lu Z-x. Antifungal activity and mechanism of fengycin in the presence and absence of commercial surfactin against *Rhizopus stolonifer*. *Journal of Microbiology*. 2011;**49**(1):146-150. DOI: 10.1007/S12275-011-0171-9
- [24] Zhao H, Shao D, Jiang C, Shi J, Li Q, Huang Q, et al. Biological activity of lipopeptides from *Bacillus*. *Applied Microbiology and Biotechnology*. 2017;**101**(15):5951-5960. DOI: 10.1007/S00253-017-8396-0
- [25] Wang S, Wang Q, Zeng X, Ye Q, Huang S, Yu H, et al. Use of the antimicrobial peptide sublancin with combined antibacterial and immunomodulatory activities to protect against methicillin-resistant *S. aureus* infection in mice. *Journal of Agricultural and Food Chemistry*. 2017;**65**(39):8595-8605. DOI: 10.1021/ACS.JAFC.7B02592
- [26] Wu C, Biswas S, Garcia De Gonzalo CV, van der Donk WA. Investigations into the mechanism of action of sublancin. *ACS Infectious Diseases*. 2019;**5**(3):454-459. DOI: 10.1021/ACSINFECDIS.8B00320/
- [27] Kang SJ, Jun JS, Moon JA, Hong KW. Surface display of p75, a *L. rhamnosus* GG derived protein, on *B. subtilis* spores and its antibacterial activity against *L. monocytogenes*. *AMB Express*. 2020;**10**(1):1-9. DOI: 10.1186/S13568-020-01073-9
- [28] Teo AYL, Tan HM. Effect of *B. subtilis* PB6 (CloSTAT) on broilers infected with a pathogenic strain of *E. coli*. *Journal of Applied Poultry Research*.

2006;**15**(2):229-235. DOI: 10.1093/JAPR/15.2.229

[29] Whelan RA, Doranalli K, Rinttilä T, Vienola K, Jurgens G, Apajalahti J. The impact of *B. subtilis* DSM 32315 on the pathology, performance, and intestinal microbiome of broiler chickens in a necrotic enteritis challenge. Poultry Science. 2019;**98**(9):3450-3463. DOI: 10.3382/PS/PEY500

[30] Ostovan R, Pourmontaseri M, Hosseinzadeh S, Shekarforoush SS. Interaction between the probiotic *B. subtilis* and *S. typhimurium* in Caco-2 cell culture. Iranian Journal of Microbiology. 2021;**13**(1):91. DOI: 10.18502/IJM.V13I1.5497

[31] Huang Q, Xu X, Mao Y-l, Huang Y, Rajput IR, Li W-f. Effects of *B. subtilis* B10 spores on viability and biological functions of murine macrophages. Animal Science Journal. 2013;**84**(3): 247-252. DOI: 10.1111/J.1740-0929.2012.01064.X

[32] Huang Q, Li YL, Xu X, Huang Y, Cui ZY, Yu DY, et al. Modulatory effects of *B. subtilis* BS02 on viability and immune responses of RAW 264.7 murine macrophages. Journal of Animal and Veterinary Advances. 2012;**11**(11):1934-1938

[33] Xu X, Huang Q, Mao Y, Cui Z, Li Y, Huang Y, et al. Immunomodulatory effects of *B. subtilis* (natto) B4 spores on murine macrophages. Microbiology and Immunology. 2012;**56**(12):817-824. DOI: 10.1111/J.1348-0421.2012.00508.X

[34] Lefevre M, Racedo SM, Ripert G, Housez B, Cazaubiel M, Maudet C, et al. Probiotic strain *B. subtilis* CU1 stimulates immune system of elderly during common infectious disease period: A randomized, double-blind

placebo-controlled study. Immunity & Ageing. 2015;**12**(1):1-11. DOI: 10.1186/S12979-015-0051-Y/

[35] Keerthi T. Immunomodulatory effect of probiotic strain *B. subtilis* MBTU PBBM1 spores in Balb/c Mice. International Journal of Engineering and Technical Research. 2014;**2**:258

[36] Bai K, Huang Q, Zhang J, He J, Zhang L, Wang T. Supplemental effects of probiotic *B. subtilis* fmbJ on growth performance, antioxidant capacity, and meat quality of broiler chickens. Poultry Science. 2017;**96**(1):74-82. DOI: 10.3382/PS/PEW246

[37] Li R, Liu J, Liu Y, Cao L, Qiu W, Qin M. Probiotic effects of *B. subtilis* on growth performance and intestinal microecological balance of growing-to-finishing pigs. Journal of Food Biochemistry. 2023;**1**:7150917. DOI: 10.1155/2023/7150917

[38] Rhee KJ, Sethupathi P, Driks A, Lanning DK, Knight KL. Role of commensal bacteria in development of gut-associated lymphoid tissues and preimmune antibody repertoire. The Journal of Immunology;**172**(2):1118-1124. DOI: 10.4049/jimmunol.172.2.1118

[39] Patch C, Pearce AJ, Cheng M, Boyapati R, Brenna JT. *B. subtilis* (BG01-4TM) improves self-reported symptoms for constipation, indigestion, and dyspepsia: A phase 1/2A randomized controlled trial. Nutrients. 2023;**15**(21):4490. DOI: 10.3390/NU15214490

[40] Cuentas AM, Deaton J, Davidson J, Ardita C. The effect of *B. subtilis* DE111 on the daily bowel movement profile for people with occasional gastrointestinal irregularity. Journal of Probiotics and Health. 2017;**5**(4):1000189. DOI: 10.4172/2329-8901.1000189

- [41] Hatanaka M, Yamamoto K, Suzuki N, Iio S, Takara T, Morita H, et al. Effect of *B. subtilis* C-3102 on loose stools in healthy volunteers. *Beneficial Microbes*. 2018;**9**(3):357-366. DOI: 10.3920/BM2017.0103
- [42] Freedman KE, Hill JL, Wei Y, Vazquez AR, Grubb DS, Trotter RE, et al. Examining the gastrointestinal and immunomodulatory effects of the novel probiotic *B. subtilis* DE111. *International Journal of Molecular Sciences*. 2021;**22**(5):2453. DOI: 10.3390/IJMS22052453
- [43] Ding H, Zhao X, Ma C, Gao Q, Yin Y, Kong X, et al. Dietary supplementation with *B. subtilis* DSM 32315 alters the intestinal microbiota and metabolites in weaned piglets. *Journal of Applied Microbiology*. 2021;**130**(1):217-232. DOI: 10.1111/jam.14767
- [44] Tang W, Qian Y, Yu B, Zhang T, Gao J, He J, et al. Effects of *B. subtilis* DSM32315 supplementation and dietary crude protein level on performance, gut barrier function and microbiota profile in weaned piglets. *Journal of Animal Science*. 2019;**97**(5):2125-2138. DOI: 10.1093/jas/skz090
- [45] Khan I, Samson SE, Grover AK. Antioxidant supplements and gastrointestinal diseases: A critical appraisal. *Medical Principles and Practice*. 2017;**26**(3):201. DOI: 10.1159/000468988
- [46] Lourenco M, Kuritza L, Westphal P, Muniz E, Pickler L, Santin E. Effects of *B. subtilis* in the dynamics of infiltration of immunological cells in the intestinal mucosa of chickens challenged with *S. minnesota*. *International Journal of Poultry Science*. 2012;**11**:630-634. DOI: 10.3923/ijps.2012.630.634
- [47] Koli D, Kadam M, Gole M, Patil A, Hajare S, Yeskal A, et al. Efficacy of *B. subtilis* (Gallipro) supplementation in *C. perfringens* challenged necrotic enteritis of broiler chicken. *Indian Journal of Animal Research*. 2018;**52**(4):619-622. DOI: 10.18805/IJAR.B-3253
- [48] Gong Y, Li H, Li Y. Effects of *B. subtilis* on epithelial tight junctions of mice with inflammatory bowel disease. *Journal of Interferon & Cytokine Research: The Official Journal of the International Society for Interferon and Cytokine Research*. 2016;**36**(2):75-85. DOI: 10.1089/JIR.2015.0030
- [49] Fujiya M, Musch MW, Nakagawa Y, Hu S, Alverdy J, Kohgo Y, et al. The *B. subtilis* quorum-sensing molecule CSF contributes to intestinal homeostasis via OCTN2, a host cell membrane transporter. *Cell Host & Microbe*. 2007;**1**(4):299-308. DOI: 10.1016/J.CHOM.2007.05.004
- [50] Okamoto K, Fujiya M, Nata T, Ueno N, Inaba Y, Ishikawa C, et al. Competence and sporulation factor derived from *B. subtilis* improves epithelial cell injury in intestinal inflammation via immunomodulation and cytoprotection. *International Journal of Colorectal Disease*. 2012;**27**(8):1039-1046. DOI: 10.1007/S00384-012-1416-8
- [51] Ricke SC, Saengkerdsub S. *Bacillus* probiotics and biologicals for improving animal and human health: Current applications and future prospects. In: Rai, Bai, editos. *Beneficial Microbes in Fermented and Functional Foods*. Boca Raton: CRC Press; 2014. pp. 341-360. DOI: 10.1201/b17912
- [52] Ma Y, Wang W, Zhang H, Wang J, Zhang W, Gao J, et al. Supplemental *B. subtilis* DSM 32315 manipulates intestinal structure and microbial composition in broiler chickens. *Scientific Reports*. 2018;**8**(1):1-13. DOI: 10.1038/s41598-018-33762-8

- [53] Wu BQ, Zhang T, Guo LQ, Lin JF. Effects of *B. subtilis* KD1 on broiler intestinal flora. Poultry Science. 2011;**90**(11):2493-2499. DOI: 10.3382/PS.2011-01529
- [54] Guo X, Li D, Lu W, Piao X, Chen X. Screening of *Bacillus* strains as potential probiotics and subsequent confirmation of the in vivo effectiveness of *B. subtilis* MA139 in pigs. Antonie van Leeuwenhoek International Journal of General and Molecular Microbiology. 2006;**90**(2):139-146. DOI: 10.1007/s10482-006-9067-9
- [55] Ciurescu G, Dumitru M, Gheorghe A, Untea AE, Drăghici R. Effect of *B. subtilis* on growth performance, bone mineralization, and bacterial population of broilers fed with different protein sources. Poultry Science. 2020;**99**(11):5960-5971. DOI: 10.1016/J.PSJ.2020.08.075
- [56] Forte C, Acuti G, Manuali E, Casagrande Proietti P, Pavone S, Trabalza-Marinucci M, et al. Effects of two different probiotics on microflora, morphology, and morphometry of gut in organic laying hens. Poultry Science. 2016;**95**(11):2528-2535. DOI: 10.3382/PS/PEW164
- [57] Sen S, Ingale SL, Kim JS, Kim KH, Kim YW, Khong C, et al. Effect of supplementation of *B. subtilis* LS 1-2 grown on citrus-juice waste and corn-soybean meal substrate on growth performance, nutrient retention, caecal microbiology and small intestinal morphology of broilers. Asian-Australasian Journal of Animal Sciences. 2011;**24**(8):1120-1127. DOI: 10.5713/AJAS.2011.10443
- [58] Kazemi SA, Ahmadi H, Karimi Torshizi MA. Evaluating two multistrain probiotics on growth performance, intestinal morphology, lipid oxidation and ileal microflora in chickens. Journal of Animal Physiology and Animal Nutrition. 2019;**103**(5):1399-1407. DOI: 10.1111/jpn.13124
- [59] Li G, Tong Y, Xiao Y, Huang S, Zhao T, Xia X. Probiotic *B. subtilis* contributes to the modulation of gut microbiota and blood metabolic profile of hosts. Comparative Biochemistry and Physiology Toxicology & Pharmacology: CBP. 2023;**272**:109712. DOI: 10.1016/J.CBPC.2023.109712
- [60] Hu Y, Dun Y, Li S, Zhao S, Peng N, Liang Y. Effects of *B. subtilis* KN-42 on growth performance, diarrhea and faecal bacterial flora of weaned piglets. Asian-Australasian Journal of Animal Sciences. 2014;**27**(8):1131. DOI: 10.5713/AJAS.2013.13737
- [61] Baker AA, Davis E, Spencer JD, Moser R, Rehberger T. The effect of a *Bacillus*-based direct-fed microbial supplemented to sows on the gastrointestinal microbiota of their neonatal piglets. Journal of Animal Science. 2013;**91**(7):3390-3399. DOI: 10.2527/JAS.2012-5821
- [62] Wang H, Kim KP, Kim IH. Influence of *B. subtilis* GCB-13-001 on growth performance, nutrient digestibility, blood characteristics, faecal microbiota and faecal score in weanling pigs. Journal of Animal Physiology and Animal Nutrition. 2019;**103**(6):1919-1925. DOI: 10.1111/jpn.13199
- [63] Zhang Q, Li J, Cao M, Li Y, Zhuo Y, Fang Z, et al. Dietary supplementation of *B. subtilis* PB6 improves sow reproductive performance and reduces piglet birth intervals. Animal Nutrition. 2020;**6**(3):278
- [64] Lv L, Mu D, Du Y, Yan R, Jiang H. Mechanism of the immunomodulatory effect of the combination of live

Bifidobacterium, *Lactobacillus*, *Enterococcus*, and *Bacillus* on immunocompromised rats. *Frontiers in Immunology*. 2021;**12**:694344. DOI: 10.3389/FIMMU.2021.694344/

[65] Cogliati S, Clementi V, Francisco M, Crespo C, Argañaraz F, Grau R. *B. subtilis* delays neurodegeneration and behavioral impairment in the Alzheimer's disease model *Caenorhabditis elegans*. *Journal of Alzheimer's Disease*. 2020;**73**(3):1035-1052. DOI: 10.3233/JAD-190837

[66] Goya ME, Xue F, Sampedro-Torres-Quevedo C, Arnaouteli S, Riquelme-Dominguez L, Romanowski A, et al. Probiotic *B. subtilis* protects against α -synuclein aggregation in *C. elegans*. *Cell Reports*. 2020;**30**(2):367. DOI: 10.1016/J.CELREP.2019.12.078

[67] Ayala FR, Bauman C, Cogliati S, Leñini C, Bartolini M, Grau R. Microbial flora, probiotics, *B. subtilis* and the search for a long and healthy human longevity. *Microbial Cell*. 2017;**4**(4):133. DOI: 10.15698/MIC2017.04.569

[68] Trotter RE, Vazquez AR, Grubb DS, Freedman KE, Grabos LE, Jones S, et al. *B. subtilis* DE111 intake may improve blood lipids and endothelial function in healthy adults. *Beneficial Microbes*. 2020;**11**(7):621-630. DOI: 10.3920/BM2020.0039

[69] Weng Y, Yao J, Sparks S, Wang KY. Nattokinase: An oral antithrombotic agent for the prevention of cardiovascular disease. *International Journal of Molecular Sciences*. 2017;**18**(3):523. DOI: 10.3390/IJMS18030523

[70] Cardinali N, Bauman C, Rodriguez Ayala F, Grau R. Two cases of type 2 diabetes mellitus successfully treated with probiotics. *Clinical Case Reports*. 2020;**8**(12):3119-3124. DOI: 10.1002/CCR3.3354

[71] Penet C, Kramer R, Little R, Spears JL, Parker J, Iyer JK, et al. A randomized, double-blind, placebo-controlled, parallel study evaluating the efficacy of *B. subtilis* MB40 to reduce abdominal discomfort, gas, and bloating. *Alternative Therapies in Health and Medicine*. 2021;**27**(S1):146-157

[72] Hatanaka M, Kanzato H, Tsuda R, Nadaoka I, Yasue M, Hoshino T, et al. Safety evaluation of the excessive intake of *B. subtilis* C-3102 in healthy Japanese adults: A randomized, placebo-controlled, double-blind, parallel-group, comparison trial. *Toxicology Reports*. 2020;**7**:46-58. DOI: 10.1016/J.TOXREP.2019.11.009

[73] Kordowski A, Tetzlaff-Lelleck VV, Speckmann B, Loh G, Künstner A, Schulz F, et al. A nutritional supplement based on a synbiotic combination of *B. subtilis* DSM 32315 and L-alanyl-L-glutamine improves glucose metabolism in healthy prediabetic subjects—A real-life post-marketing study. *Frontiers in Nutrition*. 2022;**9**:1001419. DOI: 10.3389/FNUT.2022.1001419/

[74] Toohey JC, Townsend JR, Johnson SB, Toy AM, Vantrease WC, Bender D, et al. Effects of probiotic (*B. subtilis*) supplementation during offseason resistance training in female division I athletes. *Journal of Strength and Conditioning Research*. 2020;**34**(11):3173-3181. DOI: 10.1519/JSC.0000000000002675

[75] Hanifi A, Culpepper T, Mai V, Anand A, Ford AL, Ukhanova M, et al. Evaluation of *B. subtilis* R0179 on gastrointestinal viability and general wellness: A randomised, double-blind, placebo-controlled trial in healthy adults. *Beneficial Microbes*. 2015;**6**(1):19-27. DOI: 10.3920/BM2014.0031

[76] Garvey SM, Mah E, Blonquist TM, Kaden VN, Spears JL. The probiotic *B.*

subtilis BS50 decreases gastrointestinal symptoms in healthy adults: A randomized, double-blind, placebo-controlled trial. *Gut Microbes*. 2022;**14**(1):2122668. DOI: 10.1080/19490976.2022.2122668

[77] Paytuví-Gallart A, Sanseverino W, Winger AM. Daily intake of probiotic strain *B. subtilis* DE111 supports a healthy microbiome in children attending day-care. *Beneficial Microbes*. 2020;**11**(7):611-620. DOI: 10.3920/BM2020.0022

[78] Tom Dieck H, Schön C, Wagner T, Pankoke HC, Fluegel M, Speckmann B. A synbiotic formulation comprising *B. subtilis* DSM 32315 and L-Alanyl-L-glutamine improves intestinal butyrate levels and lipid metabolism in healthy humans. *Nutrients*. 2021;**14**(1):143. DOI: 10.3390/NU14010143/S1

[79] Takimoto T, Hatanaka M, Hoshino T, Takara T, Tanaka K, Shimizu A, et al. Effect of *B. subtilis* C-3102 on bone mineral density in healthy postmenopausal Japanese women: A randomized, placebo-controlled, double-blind clinical trial. *Bioscience of Microbiota, Food and Health*. 2018;**37**(4):87-96. DOI: 10.12938/BMFH.18-006

[80] Zocco MA, Dal Verme LZ, Cremonini F, Piscaglia AC, Nista EC, Candelli M, et al. Efficacy of *Lactobacillus* GG in maintaining remission of ulcerative colitis. *Alimentary Pharmacology and Therapeutics*. 2006;**23**(11):1567-1574. DOI: 10.1111/J.1365-2036.2006.02927.X

[81] Szajewska H, Wanke M, Patro B. Meta-analysis: The effects of *L. rhamnosus* GG supplementation for the prevention of healthcare-associated diarrhoea in children. *Alimentary Pharmacology and*

Therapeutics. 2011;**34**(9):1079-1087. DOI: 10.1111/J.1365-2036.2011.04837.X

[82] Doron S, Snyderman DR, Gorbach SL. *Lactobacillus* GG: Bacteriology and clinical applications. *Gastroenterology Clinics of North America*. 2005;**34**(3):483-498. DOI: 10.1016/J.GTC.2005.05.011

[83] Yan F, Polk DB. Probiotic bacterium prevents cytokine-induced apoptosis in intestinal epithelial cells. *The Journal of Biological Chemistry*. 2002;**277**(52):50959-50965

[84] Yan F, Cao H, Cover TL, Whitehead R, Washington MK, Polk DB. Soluble proteins produced by probiotic bacteria regulate intestinal epithelial cell survival and growth. *Gastroenterology*. 2007;**132**(2):562-575. DOI: 10.1053/J.GASTRO.2006.11.022

[85] Mattila-Sandholm T, Myllärinen P, Crittenden R, Mogensen G, Fondén R, Saarela M. Technological challenges for future probiotic foods. *International Dairy Journal*. 2002;**12**(2-3):173-182. DOI: 10.1016/S0958-6946(01)00099-1

[86] Marino M, Innocente N, Calligaris S, Maifreni M, Marangone A, Nicoli MC. Viability of probiotic *L. rhamnosus* in structured emulsions containing saturated monoglycerides. *Journal of Functional Foods*. 2017;**35**:51-59. DOI: 10.1016/J.JFF.2017.05.012

[87] Cutting SM, Hong HA, Baccigalupi L, Ricca E. Oral vaccine delivery by recombinant spore probiotics. *International Reviews of Immunology*. 2009;**28**(6):487-505. DOI: 10.3109/08830180903215605

[88] Zhou Z, Gong S, Yang Y, Guan R, Zhou S, Yao S, et al. Expression of *H. pylori* urease B on the surface of *B. subtilis* spores. *Journal of Medical*

Microbiology. 2015;**64**(Pt 1):104-110.
 DOI: 10.1099/JMM.0.076430-0

[89] Ciabattini A, Parigi R, Isticato R, Oggioni MR, Pozzi G. Oral priming of mice by recombinant spores of *B. subtilis*. Vaccine. 2004;**22**(31-32):4139-4143.
 DOI: 10.1016/J.VACCINE.2004.05.001

[90] Zhao G, Miao Y, Guo Y, Qiu H, Sun S, Kou Z, et al. Development of a heat-stable and orally delivered recombinant M2e-expressing *B. subtilis* spore-based influenza vaccine. Human Vaccines & Immunotherapeutics. 2014;**10**(12):3649.
 DOI: 10.4161/HV.36122

[91] Chan BCL, Li P, Tsang MSM, Sung JCC, Kwong K WY, Zheng T, et al. Creating a vaccine-like supplement against respiratory infection using recombinant *B. subtilis* spores expressing SARS-CoV-2 Spike Protein with natural products. Molecules. 2023;**28**(13):4996.
 DOI: 10.3390/MOLECULES28134996

[92] Malaguarnera G, Giordano M, Nunnari G, Bertino G, Malaguarnera M. Gut microbiota in alcoholic liver disease: Pathogenetic role and therapeutic perspectives. World Journal of Gastroenterology: WJG. 2014;**20**(44):16639. DOI: 10.3748/WJG.V20.I44.16639

[93] Yang M, Hutchinson N, Ye N, Yin J, Guan M, Wang Z, et al. Engineered *B. subtilis* as oral probiotics to enhance clearance of blood lactate. bioRxiv. 2024:2023.11.30.569300.
 DOI: 10.1101/2023.11.30.569300

[94] Cai M, Wan J, Cai K, Song H, Wang Y, Sun W, et al. Understanding the contribution of lactate metabolism in cancer progress: A perspective from isomers. Cancers. 2022;**15**(1):87.
 DOI: 10.3390/CANCERS15010087

[95] Williamson DH, Lund P, Krebs HA. The redox state of

free nicotinamide-adenine dinucleotide in the cytoplasm and mitochondria of rat liver. The Biochemical Journal. 1967;**103**(2):514-527. DOI: 10.1042/BJ1030514

[96] Adeva-Andany M, López-Ojén M, Funcasta-Calderón R, Ameneiros-Rodríguez E, Donapetry-García C, Vila-Altesor M, et al. Comprehensive review on lactate metabolism in human health. Mitochondrion. 2014;**17**:76-100.
 DOI: 10.1016/J.MITO.2014.05.007

[97] Nørting TS, Nielsen PM, Qi H, Mikkelsen E, Hansen K, Schmidt NH, et al. Hyperbaric oxygen therapy reduces renal lactate production. Physiological Reports. 2017;**5**(6):e13217. DOI: 10.14814/PHY2.13217

[98] Sada N, Lee S, Katsu T, Otsuki T, Inoue T. Epilepsy treatment: Targeting LDH enzymes with a stiripentol analog to treat epilepsy. Science. 2015;**347**(6228):1362-1367. DOI: 10.1126/SCIENCE.AAA1299

[99] Pucino V, Cucchi D, Mauro C. Lactate transporters as therapeutic targets in cancer and inflammatory diseases. Expert Opinion on Therapeutic Targets. 2018;**22**(9):735-743.
 DOI: 10.1080/14728222.2018.1511706

[100] Thornton PK. Livestock production: Recent trends, future prospects. Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences. 2010;**365**(1554):2853-2867.
 DOI: 10.1098/RSTB.2010.0134

[101] Gadde U, Kim WH, Oh ST, Lillehoj HS. Alternatives to antibiotics for maximizing growth performance and feed efficiency in poultry: A review. Animal Health Research Reviews. 2017;**18**(1):26-45. DOI: 10.1017/S1466252316000207

- [102] Castanon JIR. History of the use of antibiotic as growth promoters in European poultry feeds. *Poultry Science*. 2007;**86**(11):2466-2471. DOI: 10.3382/PS.2007-00249
- [103] Imperial ICVJ, Ibane JA. Addressing the antibiotic resistance problem with probiotics: Reducing the risk of its double-edged sword effect. *Frontiers in Microbiology*. 2016;**7**:232849. DOI: 10.3389/FMICB.2016.01983/
- [104] Ben Y, Fu C, Hu M, Liu L, Wong MH, Zheng C. Human health risk assessment of antibiotic resistance associated with antibiotic residues in the environment: A review. *Environmental Research*. 2019;**169**:483-493. DOI: 10.1016/J.ENVRES.2018.11.040
- [105] Aarestrup FM. The livestock reservoir for antimicrobial resistance: A personal view on changing patterns of risks, effects of interventions and the way forward. *Philosophical Transactions of the Royal Society, B: Biological Sciences*. 2015;**370**(1670):20140085. DOI: 10.1098/RSTB.2014.0085
- [106] Laxminarayan R, van Boeckel T, Teillant A, Laxminarayan R, van Boeckel T. The Economic Costs of Withdrawing Antimicrobial Growth Promoters from the Livestock Sector. *OECD Food, Agriculture and Fisheries Papers*. Paris: OECD; 2015. DOI: 10.1787/5js64kst5wv1-en
- [107] Cheng G, Hao H, Xie S, Wang X, Dai M, Huang L, et al. Antibiotic alternatives: The substitution of antibiotics in animal husbandry. *Frontiers in Microbiology*. 2014;**5**:87626. DOI: 10.3389/FMICB.2014.00217/
- [108] Yuan C, Ji X, Zhang Y, Liu X, Ding L, Li J, et al. Important role of *B. subtilis* as a probiotic and vaccine carrier in animal health maintenance. *World Journal of Microbiology and Biotechnology*. 2024;**40**(9):268. DOI: 10.1007/S11274-024-04065-0
- [109] Colom J, Freitas D, Simon A, Brodkorb A, Buckley M, Deaton J, et al. Presence and germination of the probiotic *B. subtilis* DE111® in the human small intestinal tract: A randomized, crossover, double-blind, and placebo-controlled study. *Frontiers in Microbiology*. 2021;**12**:715863. DOI: 10.3389/fmicb.2021.715863
- [110] Latorre JD, Hernandez-Velasco X, Kallapura G, Menconi A, Pumford NR, Morgan MJ, et al. Evaluation of germination, distribution, and persistence of *B. subtilis* spores through the gastrointestinal tract of chickens. *Poultry Science*. 2014;**93**(7):1793-1800. DOI: 10.3382/ps.2013-03809
- [111] Saravanakumar S, Prabakaran NN, Ashokkumar R, Jamuna S. Unlocking the gut's treasure: Lipase-producing *B. subtilis* probiotic from the intestine of microstomus kitt (Lemon sole). *Applied Biochemistry and Biotechnology*. 2024;**196**(7):4273-4286. DOI: 10.1007/s12010-023-04749-7
- [112] Lee KW, Lillehoj HS, Jang SI, Lee SH. Effects of salinomycin and *B. subtilis* on growth performance and immune responses in broiler chickens. *Research in Veterinary Science*. 2014;**97**(2):304-308. DOI: 10.1016/J.RVSC.2014.07.021
- [113] Lee Y, Park I, Wickramasuriya SS, Lillehoj HS. *B. subtilis* expressing chicken NK-2 peptide enhances the efficacy of EF-1 α vaccination in eimeria maxima-challenged broiler chickens. *Animals*. 2023;**13**(8):1383. DOI: 10.3390/ANI13081383
- [114] Hong HA, Huang JM, Khaneja R, Hiep L-v, Urdaci MC, Cutting SM. The

- safety of *B. subtilis* and *B. indicus* as food probiotics. *Journal of Applied Microbiology*. 2008;**105**(2):510-520. DOI: 10.1111/J.1365-2672.2008.03773.X
- [115] Chen B, Yang Y, Wang Z, Dai X, Cao Y, Zhang M, et al. Surface display of duck hepatitis a virus type 1 VP1 protein on *B. subtilis* spores elicits specific systemic and mucosal immune responses on mice. *Probiotics and Antimicrobial Proteins*. 13 Jul 2024. DOI: 10.1007/S12602-024-10323-2
- [116] Li J, Yang M, Chen B, Wang Z, Cao Y, Yang Y, et al. Evaluation of the immunity responses in mice to recombinant *B. subtilis* displaying Newcastle Disease Virus HN Protein truncations. *Microorganisms*. 2024;**12**(3):439. DOI: 10.3390/MICROORGANISMS12030439/S1
- [117] Guo K, Huang A, Lee O. Engineering of a novel anti-dioxin *B. subtilis* probiotic for poultry feed. *Undergraduate Research in Natural and Clinical Science and Technology Journal*. 2020;**4**(1-11):1-8. DOI: 10.26685/URNCS190
- [118] Merino-Guzmán R, Latorre JD, Delgado R, Hernandez-Velasco X, Wolfenden AD, Teague KD, et al. Comparison of total immunoglobulin A levels in different samples in Leghorn and broiler chickens. *Asian Pacific Journal of Tropical Biomedicine*. 2017;**7**(2):116-120. DOI: 10.1016/J.APJT.2016.11.021
- [119] Bahaddad SA, Almalki MHK, Alghamdi OA, Sohrab SS, Yasir M, Azhar EI, et al. *Bacillus* species as direct-fed microbial antibiotic alternatives for monogastric production. *Probiotics and Antimicrobial Proteins*. 2023;**15**(1):1-16. DOI: 10.1007/S12602-022-09909-5
- [120] Xu J, Zhong F, Zhang Y, Zhang J, Huo S, Lin H, et al. Construction of *B. subtilis* strain engineered for expression of porcine β -defensin-2/cecropin P1 fusion antimicrobial peptides and its growth-promoting effect and antimicrobial activity. *Asian-Australasian Journal of Animal Sciences*. 2017;**30**(4):576. DOI: 10.5713/AJAS.16.0207
- [121] Jung K, Saif LJ. Goblet cell depletion in small intestinal villous and crypt epithelium of conventional nursing and weaned pigs infected with porcine epidemic diarrhea virus. *Research in Veterinary Science*. 2017;**110**:12-15. DOI: 10.1016/J.RVSC.2016.10.009
- [122] Gallien S, Moro A, Lediguerher G, Catinot V, Paboeuf F, Bigault L, et al. Evidence of porcine epidemic diarrhea virus (PEDV) shedding in semen from infected specific pathogen-free boars. *Veterinary Research*. 2018;**49**(1):1-9. DOI: 10.1186/S13567-018-0505-2/
- [123] Niederwerder MC, Hesse RA. Swine enteric coronavirus disease: A review of 4 years with porcine epidemic diarrhoea virus and porcine deltacoronavirus in the United States and Canada. *Transboundary and Emerging Diseases*. 2018;**65**(3):660. DOI: 10.1111/TBED.12823
- [124] Crawford K, Lager KM, Kulshreshtha V, Miller LC, Faaberg KS. Status of vaccines for porcine epidemic diarrhea virus in the United States and Canada. *Virus Research*. 2016;**226**:108-116. DOI: 10.1016/J.VIRUSRES.2016.08.005
- [125] Yu M, Wang L, Ma S, Wang X, Wang Y, Xiao Y, et al. Immunogenicity of eGFP-marked recombinant *L. casei* against transmissible gastroenteritis virus and porcine epidemic diarrhea virus. *Viruses*. 2017;**9**(10):274. DOI: 10.3390/V9100274
- [126] Liu P, Zhao J, Guo P, Lu W, Geng Z, Levesque CL, et al. Dietary corn bran

- fermented by *B. subtilis* MA139 decreased gut cellulolytic bacteria and microbiota diversity in finishing pigs. *Frontiers in Cellular and Infection Microbiology*. 2017;7:300118. DOI: 10.3389/FCIMB.2017.00526/
- [127] Wang J, Huang L, Mou C, Zhang E, Wang Y, Cao Y, et al. Mucosal immune responses induced by oral administration recombinant *B. subtilis* expressing the COE antigen of PEDV in newborn piglets. *Bioscience Reports*. 2019;39(3):BSR20182028. DOI: 10.1186/S12934-019-1151-8/
- [128] Bjerre K, Cantor MD, Nørgaard J-v, Poulsen HD, Blaabjerg K, Canibe N, et al. Development of *B. subtilis* mutants to produce tryptophan in pigs. *Biotechnology Letters*. 2017;39(2):289-295. DOI: 10.1007/S10529-016-2245-6
- [129] Peng H, Wang JQ, Kang HY, Dong SH, Sun P, Bu DP, et al. Effect of feeding *B. subtilis* natto fermentation product on milk production and composition, blood metabolites and rumen fermentation in early lactation dairy cows. *Journal of Animal Physiology and Animal Nutrition*. 2012;96(3):506-512. DOI: 10.1111/J.1439-0396.2011.01173.X
- [130] Xu H, Huang W, Hou Q, Kwok L-y, Sun Z, Ma H, et al. The effects of probiotics administration on the milk production, milk components and fecal bacteria microbiota of dairy cows. *Science Bulletin*. 2017;62(11):767-774. DOI: 10.1016/J.SCIB.2017.04.019
- [131] Sun P, Wang JQ, Deng LF. Effects of *B. subtilis* natto on milk production, rumen fermentation and ruminal microbiome of dairy cows. *Animal*. 2013;7(2):216-222. DOI: 10.1017/S1751731112001188
- [132] Al Mamun MA, Sandeman M, Rayment P, Brook-Carter P, Scholes E, Kasinadhuni N, et al. Variation in gut bacterial composition is associated with *H. contortus* parasite infection of sheep. *Animal Microbiome*. 2020;2(1):1-14. DOI: 10.1186/S42523-020-0021-3/
- [133] Xiang H, Fang Y, Tan Z, Zhong R. *Haemonchus contortus* infection alters gastrointestinal microbial community composition, protein digestion and amino acid allocations in lambs. *Frontiers in Microbiology*. 2022;12:797746. DOI: 10.3389/FMICB.2021.797746/
- [134] Yang Y, Zhang G, Wu J, Chen X, Tong D, Yang Y, et al. Recombinant HcGAPDH protein expressed on probiotic *B. subtilis* spores protects sheep from *H. contortus* infection by inducing both humoral and cell-mediated responses. *mSystems*. 2020;5(3):e00112-20. DOI: 10.1128/MSYSTEMS.00239-20
- [135] Yáñez-Mendizábal V, Zerrouh H, Viñas I, Torres R, Usall J, de Vicente A, et al. Biological control of peach brown rot (*Monilinia* spp.) by *B. subtilis* CPA-8 is based on production of fengycin-like lipopeptides. *European Journal of Plant Pathology*. 2012;132(4):609-619. DOI: 10.1007/S10658-011-9905-0/
- [136] Maksimov I-v, Singh BP, Cherepanova EA, Burkhanova GF, Khairullin RM. Prospects and applications of lipopeptide-producing bacteria for plant protection (review). *Applied Biochemistry and Microbiology*. 2020;56(1):15-28. DOI: 10.1134/S0003683820010135
- [137] Mahapatra S, Yadav R, Ramakrishna W. *B. subtilis* impact on plant growth, soil health and environment: Dr. Jekyll and Mr. Hyde. *Journal of Applied Microbiology*. 2022;132(5):3543-3562. DOI: 10.1111/JAM.15480

- [138] Qiu F, Liu W, Chen L, Wang Y, Ma Y, Lyu Q, et al. *B. subtilis* biofertilizer application reduces chemical fertilization and improves fruit quality in fertigated Tarocco blood orange groves. *Scientia Horticulturae*. 2021;281:110004. DOI: 10.1016/J.SCIENTA.2021.110004
- [139] Rostami A, Hinc K, Goshadrou F, Shali A, Bayat M, Hassanzadeh M, et al. Display of *B. pumilus* chitinase on the surface of *B. subtilis* spore as a potential biopesticide. *Pesticide Biochemistry and Physiology*. 2017;140:17-23. DOI: 10.1016/J.PESTBP.2017.05.008
- [140] Talley NJ. Functional gastrointestinal disorders as a public health problem. *Neurogastroenterology and Motility*. 2008;20(Suppl. 1):121-129. DOI: 10.1111/j.1365-2982.2008.01097.x
- [141] Ouwehand AC, Forssten S, Hibberd AA, Lyra A, Stahl B. Probiotic approach to prevent antibiotic resistance. *Annals of Medicine*. 2016;48(4):246-255. DOI: 10.3109/07853890.2016.1161232
- [142] Ishaque SM, Khosruzzaman SM, Ahmed DS, Sah MP. A randomized placebo-controlled clinical trial of a multi-strain probiotic formulation (Bio-Kult®) in the management of diarrhea-predominant irritable bowel syndrome. *BMC Gastroenterology*. 2018;18(1):71. DOI: 10.1186/s12876-018-0788-9
- [143] Moskovicz V, Ben-El R, Horev G, Mizrahi B. Skin microbiota dynamics following *B. subtilis* formulation challenge: An in vivo study in mice. *BMC Microbiology*. 2021;21(1):231. DOI: 10.1186/s12866-021-02295-y
- [144] Ben David N, Mafi M, Nyska A, Gross A, Greiner A, Mizrahi B. *B. subtilis* in PVA microparticles for treating open wounds. *ACS Omega*. 2021;6(21):13647-13653. DOI: 10.1021/acsomega.1c00790
- [145] Sarkar A, Lehto SM, Harty S, Dinan TG, Cryan JF, Burnet PWJ. Psychobiotics and the manipulation of bacteria–gut–brain signals. *Trends in Neurosciences*. 2016;39(11):763. DOI: 10.1016/j.tins.2016.09.002
- [146] Carding S, Verbeke K, Vipond DT, Corfe BM, Owen LJ. Dysbiosis of the gut microbiota in disease. *Microbial Ecology in Health and Disease*. 2015;26:26191. DOI: 10.3402/mehd.v26.26191
- [147] Luang-In V, Katisart T, Konsue A, Nudmamud-Thanoi S, Narbad A, Saengha W, et al. Psychobiotic effects of multi-strain probiotics originated from Thai fermented foods in a rat model. *Food Science of Animal Resources*. 2020;40(6):1014. DOI: 10.5851/kosfa.2020.e72
- [148] Cheng HW, Jiang S, Hu J. Gut-Brain Axis: Probiotic, *Bacillus subtilis*, Prevents Aggression via the Modification of the Central Serotonergic System [Internet]. *Oral Health by Using Probiotic Products*. London: IntechOpen; 2019. DOI: 10.5772/intechopen.86775
- [149] Jiang S, Hu JY, Cheng HW. The impact of probiotic *B. subtilis* on injurious behavior in laying hens. *Animals*. 2022;12(7):870. DOI: 10.3390/ani12070870
- [150] Wang H, Huang J, Sun L, Xu F, Zhang W, Zhan J. An efficient process for co-production of γ -aminobutyric acid and probiotic *B. subtilis* cells. *Food Science and Biotechnology*. 2019;28(1):155-163. DOI: 10.1007/s10068-018-0461-7
- [151] Díez-Gutiérrez L, San Vicente L, Luis LJ, Villarán M d C, Chávarri M. Gamma-aminobutyric acid and probiotics: Multiple health benefits and their future in the global functional food and nutraceuticals market. *Journal*

of Functional Foods. 2020;**64**:103669. DOI: 10.1016/j.jff.2019.103669

[152] Gusarov I, Gautier L, Smolentseva O, Shamovsky I, Eremina S, Mironov A, et al. Bacterial nitric oxide extends the lifespan of *C. elegans*. Cell. 2013;**152**(4):818-830. DOI: 10.1016/j.cell.2012.12.043

[153] Arnaouteli S, Bamford NC, Stanley-Wall NR, Kovács ÁT. *B. subtilis* biofilm formation and social interactions. Nature Reviews. Microbiology. 2021;**19**(9):600-614. DOI: 10.1038/s41579-021-00540-9

[154] Donato V, Ayala FR, Cogliati S, Bauman C, Costa JG, Leñini C, et al. *B. subtilis* biofilm extends *C. elegans* longevity through downregulation of the insulin-like signalling pathway. Nature Communications. 2017;**8**:14332. DOI: 10.1038/ncomms14332

[155] Dhyani P, Goyal C, Dhull SB, Chauhan AK, Singh Saharan B, Harshita, et al. Psychobiotics for mitigation of neuro-degenerative diseases: Recent advancements. Molecular Nutrition & Food Research. 2024;**68**(13):e2300461. DOI: 10.1002/mnfr.202300461

[156] Khodavirdipour A, Chamanrokh P, Alikhani MY, Alikhani MS. Potential of *B. subtilis* against SARS-CoV-2—A sustainable drug development perspective. Frontiers in Microbiology. 2022;**13**:718786. DOI: 10.3389/fmicb.2022.718786

[157] Pinu FR, Beale DJ, Paten AM, Kouremenos K, Swarup S, Schirra HJ, et al. Systems biology and multi-omics integration: Viewpoints from the metabolomics research community. Metabolites. 2019;**9**(4):76. DOI: 10.3390/metabo9040076

[158] Mugwanda K, Hamese S, van Zyl WF, Prinsloo E, Plessis M d, Dicks LMT, et al. Recent advances in genetic tools for engineering probiotic lactic acid bacteria. Bioscience Reports. 2023;**43**(1):20211299. DOI: 10.1042/BSR20211299

[159] Appala, Naidu B, Kannan K, Santhosh Kumar DP, Oliver JWK, Abbott ZD. Lyophilized *B. subtilis* ZB183 spores: 90-day repeat dose oral (gavage) toxicity study in wistar rats. Hindawi Journal of Toxicology. 2019:3042108, p. 9. DOI: 10.1155/2019/3042108

[160] The Science Behind Pre-Alcohol: A Probiotic that Digests a Byproduct of Alcohol-Built with Genetic Engineering by a Team of Microbiologists [Internet]. Available from: <https://zbiotics.com/pages/how-it-works> [Accessed: September 25, 2024]

[161] Vijayakumar KN, Bamne SS, Crasta J, Kumar CS, Bhat RM, Shuster BM, et al. *B. subtilis* ZB423: 90-day repeat dose oral (gavage) toxicity study in wistar rats. Journal of Applied Toxicology. 2024;**44**(12):1874-1885. DOI: 10.1002/jat.4677

[162] The Science Behind Sugar-to-Fiber: A Probiotic that Turns Sugar into Fiber-Built with Genetic Engineering by a Team of Microbiologists [Internet]. Available from: <https://zbiotics.com/pages/how-sugar-to-fiber-works> [Accessed: September 25, 2024]